

New Novartis study supports Entresto as foundational HFrEF therapy and in-hospital initiation in appropriate stabilized heart failure patients

Nov 11, 2018

- Entresto® (sacubitril/valsartan) outperformed commonly used heart failure medicine enalapril in landmark study; delivered significantly greater and more rapid reductions in an established biomarker for heart failure severity and prognosis(1)
- Significant 46% reduction in serious clinical outcomes endpoint, primarily by reducing death and heart failure re-hospitalization, compared to enalapril over 8 weeks in pre-specified exploratory analysis(1)
- Safety and tolerability comparable to enalapril, including rates of hypotension, hyperkalemia, renal complications and risk of angioedema(1)
- Currently, patients with heart failure with reduced ejection fraction (HFrEF) who need to be admitted to the hospital for an acute decompensation event are at high risk of short-term death and costly heart failure re-hospitalizations(2,3)

EAST HANOVER, N.J., Nov. 11, 2018 /PRNewswire/ -- Novartis announced today results of the landmark PIONEER-HF trial showing that in-hospital initiation of Entresto tablets provided superior benefit compared to enalapril—a heart failure medication commonly used—in patients with HFrEF who had been stabilized following admission for an acute decompensation heart failure (ADHF) event.¹ Results were presented as a late-breaker at the American Heart Association's Scientific Sessions 2018 and published in The New England Journal of Medicine.

Patients on Entresto in PIONEER-HF had a 29% greater reduction in time-averaged N-terminal pro-B-type natriuretic peptide (NT-proBNP) at weeks 4 and 8 (the primary endpoint) compared to enalapril patients (95% CI: 0.63, 0.81; $P < 0.0001$).¹ Significant reductions in NT-proBNP were observed in Entresto patients as early as 1 week after treatment initiation.¹ NT-proBNP is an established biomarker used to assess the severity and determine the prognosis of heart failure.⁴ Notably, the superior NT-proBNP reduction with Entresto was consistent across diverse HFrEF patient populations stabilized following admission for ADHF, including those newly diagnosed with HFrEF, those not receiving an ACEi/ARB treatment and African Americans.¹

In a pre-specified exploratory analysis of PIONEER-HF, Entresto also showed a significant 46% reduction in the risk of a composite of death, heart failure re-hospitalization (hospital stay >24 hours), requirement for left ventricular assist device (LVAD) insertion or listing for cardiac transplantation compared to enalapril over 8 weeks.¹ This result was driven primarily by reductions in death and HF re-hospitalization among patients treated with Entresto.¹ There were no new safety signals identified.¹ Entresto is indicated to reduce the risk of CV death or HF hospitalization in patients with chronic heart failure (NYHA Class II-IV) and reduced ejection fraction.⁵

"The results of this landmark study should help inform our basic approach to treating hospitalized patients with

acute heart failure," said Eric Velazquez, MD, Berliner Professor of Cardiology at Yale School of Medicine and PIONEER-HF principal study investigator. "With the PIONEER-HF trial results—once acute heart failure is diagnosed, patients are hemodynamically stabilized, and a low ejection fraction is confirmed—sacubitril/valsartan should be started promptly to reduce neurohormonal activation and reduce the risk of post-discharge heart failure hospitalization."

"PIONEER-HF further confirmed the safety and benefit, as reflected in the reduction of an important biomarker, of in-hospital initiation of Entresto treatment in HFrEF patients stabilized following ADHF," said Shreeram Aradhye, MD, Chief Medical Officer and Global Head, Medical Affairs, Novartis Pharmaceuticals. "Together with data from the PARADIGM-HF study, which demonstrated the superior benefit of Entresto compared to an ACEi on cardiovascular mortality and HF hospitalizations in ambulatory patients, there is now consistent evidence in the in and outpatient settings supporting the use of Entresto. Through the Entresto scientific program, we are reimagining the standard of care for HFrEF patients and the use of Entresto as foundation therapy."

Patients with heart failure who need to be admitted to the hospital due to decompensation of their disease are at high short-term risk of death or being re-admitted to the hospital.^{2,3} PIONEER-HF showed that Entresto can be initiated in the hospital in these patients safely, once they have been stabilized, with a tolerability profile comparable to enalapril.¹ Rates of hypotension (low blood pressure), hyperkalemia (elevated levels of potassium) or renal (kidney) complications were similar and there was no greater risk of angioedema in patients treated with Entresto.¹ There were no new safety signals identified, and the safety profile was comparable to that seen in PARADIGM-HF.^{1,6} These results add to learnings from the TRANSITION study (an open label parallel group design), which explored the initiation of Entresto shortly after patients were stabilized following an acute heart failure episode, both in the hospital and out-patient settings.^{7,8} Initial TRANSITION results were presented at the European Society of Cardiology Congress in August 2018.⁷ Additional biomarker data from TRANSITION will be presented at AHA.

About the PIONEER-HF Trial

PIONEER-HF is a prospective, multicenter, double-blind, randomized, controlled trial designed to assess the safety, tolerability, and efficacy of in-hospital initiation of Entresto compared with enalapril in appropriate, stable HFrEF patients who had been admitted for acute decompensation.^{1,9}

The study enrolled patients 18 years of age and older with ejection fraction (EF) $\leq 40\%$ and an elevated amino terminal-pro b-type natriuretic peptide (NT-proBNP) ≥ 1600 pg/mL or b-type natriuretic peptide (BNP) ≥ 400 pg/mL, irrespective of both duration of diagnosis or treatment with angiotensin converting-enzyme inhibitors (ACEi) and/or angiotensin receptor blockers (ARB).^{1,9}

A total of 881 patients, with an average age of 61 years, were randomly assigned to in-hospital initiation of Entresto or enalapril twice daily, following stabilization.^{1,9} Patients were eligible for screening no earlier than 24 hours and up to 10 days from initial presentation while still hospitalized.^{1,9} All patients were treated with the objective to optimize therapy to the maximum tolerated dose based on an SBP-based algorithm.^{1,9} Patients were predominantly male (approximately 72%), and half of patients had a BMI > 30 kg/m².^{1,9} Notably, more than one-third of patients (36%) were African American.^{1,9} Approximately 34% of patients were newly diagnosed, having no prior history of heart failure, and slightly more than 50% of patients were not receiving ACEi/ARB therapy at the time of admission.^{1,9}

The primary endpoint was the time-averaged proportional change in NT-proBNP from baseline through weeks 4 and 8.^{1,9} Patients treated with Entresto achieved a 47% reduction from baseline in time-averaged NT-

proBNP, compared to a 25% reduction with enalapril, translating into a statistically significant 29% greater reduction with Entresto over the ACE inhibitor (95% CI: 0.63, 0.81; $P < 0.0001$).¹ Significant reductions in NT-proBNP were observed in Entresto patients as early as one week after treatment initiation.¹ Rates of serious adverse events occurring with a frequency of $\geq 0.5\%$ were similar between the Entresto and enalapril groups.¹

Safety and tolerability analyses found:

- Similar levels of symptomatic hypotension in both arms (enalapril 12.7%, Entresto 15.0%; RR: 1.18; 95% CI: 0.85, 1.64). Low BP-related AEs were similar between the groups.¹
- Hyperkalemia was comparable between the two groups (enalapril 9.3% vs. Entresto 11.6% RR: 1.25; 95% CI: 0.84, 1.84).¹
- Kidney-related AEs between the groups were similar (enalapril $n = 75$, Entresto $n = 75$, RR: 1.00; CI: 0.75, 1.34).¹
- There were 6 adjudicated cases of angioedema in the enalapril arm (all in African Americans) versus 1 with Entresto (in a Caucasian patient).¹
- Nearly 60% of patients were able to be up-titrated to the highest approved dose of Entresto within 6 weeks.¹

In a pre-specified exploratory analysis, the composite endpoint of death, re-hospitalization for HF, LVAD implantation or listing for cardiac transplant occurred in 41 (9.3%) patients in the Entresto group and 74 (16.8%) in the enalapril group (hazard ratio [HR] 0.54, 95% CI 0.37–0.79; $p = .001$).¹ The benefit was driven by reductions in death and re-hospitalization among patients treated with Entresto. The number needed to treat to prevent one such clinical event during 8 weeks of follow up was 13.¹

About NT-proBNP

NT-proBNP is a biomarker commonly used to assess the severity and determine the prognosis of heart failure.⁴ Levels of NT-proBNP increase when heart muscle cells are subjected to stress (such as stretching) that occurs in people with heart failure.⁴ Studies suggest that HF patients with elevated NT-proBNP are at an increased risk of CV death or HF hospitalization and that reducing levels of NT-proBNP in people with heart failure is associated with a lower risk of these negative clinical outcomes.⁴ Entresto was also shown to reduce plasma NT-proBNP compared with enalapril in the PIONEER-HF and PARADIGM-HF trials.^{1,6}

About Heart Failure

Heart failure (HF) is a chronic and progressive condition, which impacts 6.5 million Americans and is the leading cause of hospitalization among Americans over the age of 65.^{10,11} About half of people with HF have heart failure with reduced ejection fraction (HFrEF), also known as systolic HF.^{12,13} Reduced ejection fraction means the heart does not contract with enough force, so less blood is pumped out.¹⁴ HF presents a major and growing health-economic burden that currently exceeds \$30 billion in the United States, which accounts for both direct and indirect costs.¹⁵

Novartis has established the largest global clinical program in the HF disease area across the pharma industry to date. Known as FortiHFy, it is comprised of more than 40 active or planned clinical studies designed to generate an array of additional data on symptom reduction, efficacy, quality of life benefits and real-world evidence with Entresto, as well as to extend understanding of heart failure.

About Entresto

Entresto is a prescription medicine used to reduce the risk of death and hospitalization in people with certain types of long-lasting (chronic) heart failure.⁵ Entresto is usually used with other heart failure therapies, in place

of an ACE inhibitor or other ARB therapy.⁵ Entresto is a twice-a-day prescription medicine that reduces the strain on the failing heart. It does this by enhancing the beneficial neurohormonal systems (natriuretic peptide system) while simultaneously inhibiting the harmful effects of the overactive renin-angiotensin-aldosterone system (RAAS).^{5,16} Most other heart failure medicines only block the harmful effects of the overactive RAAS. Entresto contains the neprilysin inhibitor sacubitril and the angiotensin receptor blocker (ARB) valsartan.¹² Entresto film-coated tablets are available in three dosage strengths: 24/26 mg, 49/51 mg, and 97/103 mg (sacubitril/valsartan).⁵ These doses are referred to as 50 mg, 100 mg, and 200 mg in the clinical trial literature including The New England Journal of Medicine publication of the results of PARADIGM-HF.⁵ The target maintenance dose of Entresto is 97/103 mg twice daily.⁵

IMPORTANT SAFETY INFORMATION

Entresto can harm or cause death to an unborn baby. Patients should talk to their doctor about other ways to treat heart failure if they plan to become pregnant. If a patient gets pregnant while taking Entresto, she should tell her doctor right away.

Patients are not to take Entresto if they are allergic to sacubitril or valsartan or any of the ingredients in Entresto; have had an allergic reaction including swelling of the face, lips, tongue, throat or trouble breathing while taking a type of medicine called angiotensin-converting enzyme (ACE) inhibitor or angiotensin II receptor blocker (ARB); or take an ACE inhibitor medicine. Patients are not to take Entresto for at least 36 hours before or after they take an ACE inhibitor medicine. Patients should talk with their doctor or pharmacist before taking Entresto if they are not sure if they take an ACE inhibitor medicine. Patients are not to take Entresto if they have diabetes and take a medicine that contains aliskiren.

Before they take Entresto, patients should tell their doctor about all of their medical conditions, including if they have kidney or liver problems; or a history of hereditary angioedema; are pregnant or plan to become pregnant; are breastfeeding or plan to breastfeed. Patients should either take Entresto or breastfeed. They should not do both.

Patients should tell their doctor about all the medicines they take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. They should especially tell their doctor if they take potassium supplements or a salt substitute; nonsteroidal anti-inflammatory drugs (NSAIDs); lithium; or other medicines for high blood pressure or heart problems such as an ACE inhibitor, ARB, or aliskiren.

Entresto may cause serious side effects including serious allergic reactions causing swelling of the face, lips, tongue, and throat (angioedema) that may cause trouble breathing and death. Patients are to get emergency medical help right away if they have symptoms of angioedema or trouble breathing. Patients are not to take Entresto again if they have had angioedema while taking Entresto. People who are black or who have had angioedema may have a higher risk of having angioedema if they take Entresto. Entresto may cause low blood pressure (hypotension). Patients are to call their doctor if they become dizzy or lightheaded, or they develop extreme fatigue. Entresto may cause kidney problems or an increased amount of potassium in the blood.

The most common side effects were low blood pressure, high potassium, cough, dizziness, and kidney problems.

Please see full Prescribing Information, including Boxed WARNING available at <http://www.pharma.us.novartis.com/product/pi/pdf/entresto.pdf>.

Patients are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Novartis is committed to providing patients with affordable access and resources through Entresto Central. For more information, please call 1-888-ENTRESTO or visit www.entresto.com.

Disclaimer

This press release contains forward-looking statements within the meaning of the United States Private Securities Litigation Reform Act of 1995. Forward-looking statements can generally be identified by words such as "potential," "can," "will," "plan," "expect," "anticipate," "look forward," "believe," "committed," "investigational," "pipeline," "launch," or similar terms, or by express or implied discussions regarding potential marketing approvals, new indications or labeling for the investigational or approved products described in this press release, or regarding potential future revenues from such products. You should not place undue reliance on these statements. Such forward-looking statements are based on our current beliefs and expectations regarding future events, and are subject to significant known and unknown risks and uncertainties. Should one or more of these risks or uncertainties materialize, or should underlying assumptions prove incorrect, actual results may vary materially from those set forth in the forward-looking statements. There can be no guarantee that the investigational or approved products described in this press release will be submitted or approved for sale or for any additional indications or labeling in any market, or at any particular time. Nor can there be any guarantee that such products will be commercially successful in the future. In particular, our expectations regarding such products could be affected by, among other things, the uncertainties inherent in research and development, including clinical trial results and additional analysis of existing clinical data; regulatory actions or delays or government regulation generally; global trends toward health care cost containment, including government, payor and general public pricing and reimbursement pressures; our ability to obtain or maintain proprietary intellectual property protection; the particular prescribing preferences of physicians and patients; general political and economic conditions; safety, quality or manufacturing issues; potential or actual data security and data privacy breaches, or disruptions of our information technology systems, and other risks and factors referred to in Novartis AG's current Form 20-F on file with the US Securities and Exchange Commission. Novartis is providing the information in this press release as of this date and does not undertake any obligation to update any forward-looking statements contained in this press release as a result of new information, future events or otherwise.

About Novartis

Novartis is reimagining medicine to improve and extend people's lives. As a leading global medicines company, we use innovative science and digital technologies to create transformative treatments in areas of great medical need. In our quest to find new medicines, we consistently rank among the world's top companies investing in research and development. Novartis products reach nearly 1 billion people globally and we are finding innovative ways to expand access to our latest treatments. About 125,000 people of more than 140 nationalities work at Novartis around the world. Novartis Pharmaceuticals Corporation, a US affiliate of Novartis, is located in East Hanover, NJ.

Find out more at www.novartis.com

Novartis is on Twitter. Sign up to follow @Novartis at <http://twitter.com/novartis>

For Novartis multimedia content, please visit www.novartis.com/news/media-library

For questions about the site or required registration, please contact media.relations@novartis.com

References

1. Velazquez E, Morrow D, DeVore, A, et al., Angiotensin-Nepriylsin Inhibition in Acute Decompensated Heart Failure. N Engl J Med. 2018. doi: 10.1056/NEJMoa1812851.
2. Dharmarajan K, Hsieh AF, Lin Z, et al., Diagnoses and Timing of 30-Day Readmissions after

- Hospitalization For Heart Failure, Acute Myocardial Infarction, or Pneumonia. JAMA. 2013;309(4):355-363.
3. Bueno H, Ross JS, Wang Y, et al., Trends in Length of Stay and Short-Term Outcomes among Medicare Patients Hospitalized for Heart Failure: 1993-2008. JAMA. 2010;303(21):2141-2147.
 4. Ndumele C, et al. NT-proBNP and Heart Failure Risk Among Individuals With and Without Obesity: The ARIC Study. Circulation. 2016; 133:631-638.
 5. ENTRESTO [prescribing information]. East Hanover, NJ: Novartis Pharmaceuticals Corp; November 2017.
 6. McMurray JJV, Packer M, Desai AS, et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. N Engl J Med. 2014;371:993-1004. doi: 10.1056/NEJMoa1409077.
 7. Wachter R. et al., Initiation of sacubitril/valsartan in hospitalized patients with heart failure with reduced ejection fraction after hemodynamic stabilization: Primary results of the TRANSITION study. Data presented at: ESC 2018, Aug 25-29; Munich, Germany.
 8. Pascual-Figal D, et al., Rationale and design of TRANSITION: a randomized trial of pre-discharge vs. post-discharge initiation of sacubitril/valsartan. ESC Heart Fail. 2018 Apr;5(2):327-336.
 9. Velazquez EJ, et al., Rationale and design of the comparison of sacubitril/valsartan versus Enalapril on Effect on nt-pRo-bnp in patients stabilized from an acute Heart Failure episode (PIONEER-HF) trial. Am Heart J. 2018 Apr;198:145-151.
 10. Benjamin EJ, Blaha MJ, Chiuve SE, et al. e. Heart disease and stroke statistics—2017 update: a report from the American Heart Association. Circulation. 2017;135:00-00. doi: 10.1161/CIR.0000000000000485.
 11. Weir LM, Pfuntner A, Maeda J, et al. HCUP facts and figures: statistics on hospital-based care in the United States, 2009. Rockville, MD: Agency for Healthcare Research and Quality, 2011.
 12. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: A report of the American College of Cardiology Foundation/American Heart Association task force on practice guidelines. Circulation. 2013;128:e240-e327.
 13. Owan TE, Hodge DO, Herges RM, et al. Trends in prevalence and outcome of heart failure with preserved ejection fraction. N Engl J Med. 2006;355:251-259.
 14. Ejection Fraction Heart Failure Measurement. American Heart Association Website. http://www.heart.org/HEARTORG/Conditions/HeartFailure/SymptomsDiagnosisofHeartFailure/Ejection-Fraction-Heart-Failure-Measurement_UCM_306339_Article.jsp (link is external). Published March 24, 2015. Accessed November 1, 2018.
 15. Heidenreich PA, Albert NM, Allen LA, et al. Forecasting the impact of heart failure in the United States: a policy statement from the American Heart Association. Circ Heart Fail. 2013;6:606-619.
 16. Langenickel T, Dole W. Angiotensin receptor-neprilysin inhibition with LCZ696: a novel approach for the

treatment of heart failure. Drug Discov Today. 2012;4: e131-139.

Novartis Media Relations

Central media line: +41 61 324 2200

E-mail: media.relations@novartis.com

Sandra Blum

Novartis Cardio-Metabolic Communications

+41 61 324 6441 (direct)

+41 79 540 7348 (mobile)

sandra.blum@novartis.com

Eric Althoff

Novartis Global Media Relations

+41 61 324 7999 (direct)

+41 79 593 4202 (mobile)

eric.althoff@novartis.com

Phil McNamara

Novartis US Pharma Communications

+1 862 778 0218 (direct)

+1 862 274 5255 (mobile)

philip.mcnamara@novartis.com

Novartis Investor Relations

Central investor relations line: +41 61 324 7944

E-mail: investor.relations@novartis.com

Central

North America

Samir Shah

+41 61 324 7944

Richard Pulik

+1 212 830 2448

Pierre-Michel Bringer

+41 61 324 1065

Cory Twining

+1 212 830 2417

Thomas Hungerbuehler

+41 61 324 8425

Source URL: <https://prod1.novartis.com/us-en/news/media-releases/new-novartis-study-supports-entresto-foundational-hfref-therapy-and-hospital-initiation-appropriate-stabilized-heart-failure-patients>

List of links present in page

1. <https://prod1.novartis.com/us-en/us-en/news/media-releases/new-novartis-study-supports-entresto-foundational-hfref-therapy-and-hospital-initiation-appropriate-stabilized-heart-failure-patients>
2. <http://www.pharma.us.novartis.com/product/pi/pdf/entresto.pdf>
3. <http://www.fda.gov/medwatch>
4. <http://www.entresto.com/>
5. <http://www.novartis.com/>
6. <http://twitter.com/novartis>
7. <http://www.novartis.com/news/media-library>
8. <mailto:media.relations@novartis.com>
9. http://www.heart.org/HEARTORG/Conditions/HeartFailure/SymptomsDiagnosisofHeartFailure/Ejection-Fraction-Heart-Failure-Measurement_UCM_306339_Article.jsp
10. <mailto:media.relations@novartis.com>
11. <mailto:eric.althoff@novartis.com>
12. <mailto:sandra.blum@novartis.com>
13. <mailto:philip.mcnamara@novartis.com>
14. <mailto:investor.relations@novartis.com>